

2020

Social networks differ depending on aggressive and affiliative interactions of captive ring-tailed lemurs (*Lemur catta*)

Marcus, K.

Marcus, K. (2020) 'Social networks differ depending on aggressive and affiliative interactions of captive ring-tailed lemurs (*Lemur catta*)', *The Plymouth Student Scientist*, 13(1), p. 64-96.

<http://hdl.handle.net/10026.1/16506>

The Plymouth Student Scientist

University of Plymouth

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Appendix

Table A1: Total observed and expected counts of aggressive and submissive behaviours emitted during feeding sessions by each dominance rank in addition to their contribution to chi-squared. Observed counts that were significantly different from the expected count are coloured red.

Rank		Move Away	Cower	Submit Food	Displace	Lunge	Chase	Cuff	Take Food	Bite	Fight
High	Observed	110	9	5	184	102	140	192	47	5	7
	Expected	337.2	37.9	22.3	112.8	60.8	82.1	113.9	26.8	3.4	3.9
	Contribution to Chi-square	153.1	22.1	13.5	45.1	27.8	40.9	53.6	15.2	0.8	2.5
Medium	Observed	197	23	8	13	9	4	9	1	1	0
	Expected	110.3	12.4	7.3	36.9	19.9	26.8	37.3	8.8	1.1	1.3
	Contribution to Chi-square	68.2	9.1	0.1	15.5	9.7	19.4	21.4	6.9	0.01	1.3
Low	Observed	297	36	27	5	1	3	3	0	0	0
	Expected	156.6	17.6	10.4	52.4	28.3	38.1	52.9	12.4	1.6	1.8
	Contribution to Chi-square	125.9	19.2	26.7	42.8	26.3	32.3	47.1	12.4	1.6	1.8

Critical chi-squared value = 10.83

Table A2: Total observed and expected counts of aggressive and submissive behaviours emitted during non-feeding sessions by each dominance rank in addition to their contribution to chi-squared. Observed counts that were significantly different from the expected count are coloured red.

Rank		Move Away	Cower	Displace	Lunge	Chase	Cuff	Grab	Bite	Fight
High	Observed	30	2	42	19	44	57	11	3	10
	Expected	88.9	11.2	28.5	15.1	22.3	39.3	4.6	2.3	5.8
	Contribution to Chi-square	39.1	7.5	6.4	1.1	21.1	7.9	8.8	0.2	3.1
Medium	Observed	89	7	15	10	11	26	0	2	2
	Expected	66.1	8.3	21.2	11.2	16.6	29.2	3.4	1.7	4.3
	Contribution to Chi-square	7.9	0.2	1.8	0.1	1.9	0.4	3.4	0.03	1.2
Low	Observed	112	20	17	10	3	19	1	1	3
	Expected	75.9	9.5	24.3	12.8	19.1	33.5	3.9	1.9	4.9
	Contribution to Chi-square	17.2	11.5	2.2	0.6	13.5	6.3	2.2	0.5	0.8

Critical chi-squared value = 10.83