

2022-10

Mesoscale eddies and the impact of coastal iron supply on primary production in the South Pacific Subtropical Front

Chiswell, SM

<http://hdl.handle.net/10026.1/19946>

10.1016/j.dsr.2022.103852

Deep Sea Research Part I: Oceanographic Research Papers

Elsevier

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

This is a submitted draft and does not contain images as these were uploaded separately. For the most up to date version, which is correctly formatted and includes tables/figures, please use the open access link

<https://doi.org/10.1016/j.dsr.2022.103852>

1

2 **Mesoscale eddies and the impact of coastal iron supply on primary production in**
3 **the South Pacific Subtropical Front**

4 ¹Stephen M. Chiswell5 ²Philip W. Boyd6 ^{3,4}Sylvia G. Sander7 ⁵Michael J. Ellwood8 ⁶Angela Milne

9

10 ¹National Institute of Water and Atmospheric Research (NIWA), Wellington, New
11 Zealand12 ²Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS
13 7004, Australia14 ³Department of Chemistry, NIWA/University of Otago Research Centre for
15 Oceanography, University of Otago, Dunedin 9054, New Zealand16 ⁴Marine Mineral Resources, Dynamics of the Ocean Floor, GEOMAR Helmholtz
17 Centre for Ocean Research Kiel, 24148 Kiel, Germany18 ⁵Research School of Earth Sciences, Australian National University, Canberra, ACT
19 2601, Australia20 ⁶School of Geography, Earth and Environmental Sciences, University of Plymouth,
21 Plymouth PL4 8AA, United Kingdom

22

23 Corresponding author: Stephen Chiswell: s.chiswell@niwa.cri.nz

ABSTRACT

24 Subtropical and Subantarctic waters either side of the southern hemisphere
25 Subtropical Front are considered iron-limited, suggesting production within the front
26 is dependent on a supply of iron from atmospheric deposition, zonal advection of
27 coastal water, or upwelling. We present the results from a one-day biogeochemical
28 survey in Subtropical Water east of the North Island, New Zealand, in a region where
29 mesoscale cyclonic and anticyclonic eddies entrain chlorophyll in filaments around
30 the eddies. There was no significant relationship between upper-mixed layer
31 chlorophyll and any physical or macronutrient quantity. However, chlorophyll was
32 significantly positively correlated with dissolved iron. A simple model suggests that
33 while vertical entrainment of iron into the upper mixed layer occurred, most of the
34 dissolved iron in the eddy was due to entrainment of high-iron coastal water into low-
35 iron offshore Subtropical Water, and that this iron supports primary production in
36 otherwise iron-deficient water. We suggest that a significant component of the total
37 primary production within the STF may be determined by mesoscale eddy induced
38 lateral advection of iron.

39

40 *Keywords:* Iron supply; Neritic water; Subtropical water; Primary production; South-
41 west Pacific Ocean; New Zealand

42

1. Introduction

43 Primary production at subtropical latitudes in the South Pacific Ocean is
44 determined by the nutrient and physical characteristics of two main surface water
45 masses in the region. To the south, Subantarctic Water (SAW) is recognised as high
46 nitrate-low chlorophyll water (HNLC) where surface summer and winter nitrate
47 concentrations are typically 7 and 15 mmol m^{-3} , respectively (Johnson et al., 2017),
48 yet mean sea surface chlorophyll (SSC) is typically only 0.3 mg Chl m^{-3} (Banse and
49 English, 1997). SAW is considered iron-limited (Abraham et al., 2000; Boyd et al.,
50 1999; Boyd et al., 2000), and it is often concluded that overall production in SAW
51 depends on the input of dissolved iron (dFe). To the north, Subtropical Water (STW)
52 is regarded as being at least seasonally, if not year round, oligotrophic, where surface
53 nitrate concentrations are typically $<2.5 \text{ mmol m}^{-3}$ year round (Chiswell et al., 2022;
54 Ellwood et al., 2018). Although STW has not been as well studied as SAW, it is also
55 likely to be seasonally depleted in dFe both in the western (Ellwood et al., 2008) and
56 eastern (Blain et al., 2008) South Pacific Ocean.

57 These two water masses meet at the Subtropical Front (STF), which spans the
58 globe near 40°S , and is a region of high primary production, where the mean SSC can
59 exceed 0.75 mg Chl m^{-3} (Figure 1a). Since both STW and SAW are probably iron-
60 limited, primary production in the STF is likely to be supported by an influx of dFe.
61 Three commonly discussed candidates for iron enrichment in the ocean are
62 atmospheric deposition (e.g. Jickells and Moore, 2015), zonal advection of high-iron
63 neritic water (e.g. Boyd et al., 2012; Ellwood et al., 2014; Graham et al., 2015), and
64 mesoscale-eddy driven vertical pumping (e.g. Uchida et al., 2020), and it seems that
65 primary production along the STF must be controlled by some combination of these
66 three mechanisms. Since mean SSC tends to show higher values closer to land (Figure
67 1a), the two most likely mechanisms would appear to be atmospheric deposition and
68 zonal advection. However, there is no consensus on which mechanism dominates, and
69 there is not yet a complete understanding of the role of iron in STF production.

70 The mean circulation east of the North Island is dominated by the warm-core
71 anticyclonic Wairarapa Eddy, which recirculates STW from the northern edge of the
72 STF so that there is a region of enhanced SSC extending from East Cape to the
73 Chatham Islands (Figure 1b). Mean SSC is depressed at the centre of the eddy, and
74 this depressed biomass has been related to the deeper mixed layer in the eddy centre

75 limiting primary production (Bradford et al., 1982; Waite et al., 2007). However, this
76 picture is only true in the mean, and at any given time, the region is dominated by a
77 complex pattern of anticyclonic and cyclonic mesoscale eddies likely shed near the
78 East Cape and propagating south (Chiswell, 2005).

79 During September-October 2012, a two-week cruise was made east of the North
80 Island, New Zealand, to investigate the role of iron in the evolution of the spring
81 bloom in STW. The spring bloom was not spatially heterogeneous, and a satellite
82 image transmitted to the ship showed that SSC in the region was dominated by eddy
83 mixing, with filaments of high SSC entrained around mesoscale eddies. As a result of
84 this image, a one-day survey was designed to investigate an anticyclonic eddy and its
85 impact on production.

86 This article presents the results of this survey, where we found no discernible
87 differences in the water mass properties, mixed-layer depths, or macronutrient
88 concentrations between regions of high and low SSC in and around the eddy. The
89 only measurable differences were that dFe was higher in regions of high SSC and
90 where the upper mixed layer was deeper. A simple model suggests that while vertical
91 entrainment of iron into the upper mixed layer occurred, most of the dissolved iron
92 signal was due to the eddy entraining high-iron coastal water into low-iron offshore
93 Subtropical Water, and that near-surface chlorophyll was a response to this iron. The
94 resulting implication is that a significant component of the total primary production
95 within the STF is determined by mesoscale-eddy induced zonal advection of iron.

2. Data and Methods

96 The cruise was made to the study site east of the North Island, nominally at
97 180°E, 39°S (Figure 2) from mid-September until early October 2012 to study the
98 2012 spring bloom. Details of the cruise and the results, including the bloom
99 development, are given in Chiswell et al. (2019).

100 The one-day survey track was based on a MODIS image of 30 September
101 (Figure 3) showing an anticyclonic eddy centred near 180°E, 39°S (labelled A₁ in
102 Figure 2). The survey track started near the centre of the eddy as suggested by the
103 surface chlorophyll pattern, and was then made about 50 km to the north/north-west
104 before returning to the eddy centre. Data from a shipboard ADCP were used to

105 determine near-surface ocean velocity and used to guide the survey in locating the
106 eddy centre.

107 Both vertical profiles from CTD casts, and near-surface underway data from a
108 towed-fish and the ship's sea chest instrumentation were collected.

109 Twelve CTD casts were made, from 03:30 on 2 October until 04:30 on 3
110 October (New Zealand Standard Time), 4 casts (2 through 5) were made during
111 daylight, one cast (6) was made in twilight while the rest were made during the night.
112 The CTD casts were made using a standard SeaBird 911 and carousel water sampler.
113 Temperature, salinity, transmissivity, photosynthetic active radiation (PAR), and
114 fluorescence profiles were measured to at least 350 m depth. Water samples for
115 chlorophyll extractions and nutrient analyses were collected on up-casts using 24 10-L
116 Niskin bottles mounted on the CTD rosette.

117 Chlorophyll derived from the CTD fluorometer was compromised by non-
118 photochemical quenching (e.g., Carberry et al., 2019) during the daylight casts, so the
119 CTD fluorometer data are not used in the analysis. Instead, following Bishop (1999),
120 particulate organic matter (POM) was taken to be negatively proportional to the beam
121 attenuation coefficient, c , derived from the CTD transmissometer as $c = \ln(T_r) / r$,
122 where T_r is the transmissivity, and r is the path length. With no local calibration
123 between POM and c , POM was normalised to have a maximum value of 1.0. A
124 comparison of POM and CTD-derived chlorophyll (Chl) from night-time casts (to
125 avoid quenching issues) for depths $5 > 30$ m over the entire 2-week cruise indicated a
126 not quite linear relationship ($r^2 = 0.89$, Figure 4g). To the extent that Chl is an
127 indicator of biomass, this supports the assumption that POM can be taken as a proxy
128 for phytoplankton biomass.

129 The one-percent light level for each daylight cast was computed by fitting an
130 exponential decay with depth function, $I = I_{sfc} \exp(-k \times z)$ to each PAR profile, and
131 computing the one-percent light level as $Z_{100} = \ln(0.01) / k$.

132 Macronutrients from the CTD up-cast samples were determined using an
133 automated micro-segmented flow analyser with digital detection (Pickmere, 1998).
134 The standard error in the automated analyses for each nutrient was estimated as the
135 standard deviation of samples taken over a 6-hour night-time period earlier during the

136 two-week cruise when the ship was on station. These overall standard deviations were
137 0.390 , 0.064, 0.23 and 0.093 $\mu\text{mol L}^{-1}$ for nitrate, phosphate, silicate and ammonium,
138 respectively. All indicators suggested variations seen in nutrient profiles were outside
139 the error for individual measurements.

140 Near-surface dFe was measured from sea water pumped from a towed ‘trace-
141 metal fish’, and was determined using flow injection analysis (Floor et al., 2015;
142 Obata et al., 2002). Further analytical details are described in Ellwood et al. (2015).

143 Continuous near-surface measurements of temperature, salinity and
144 fluorescence were made with a thermosalinograph (Seabird 38 and 21 sensors) and
145 fluorometer (Wetlab ECO triplet) in the ship’s sea chest. The sea chest also contained
146 an inline Fast Repetition Rate Fluorometer (Chelsea Instruments FASTtracka FRRF)
147 that provided measurements every 2 minutes of near-surface minimum (F_o) and
148 maximum (F_m) fluorescence in the dark-acclimated state.

149 There was almost certainly some NPQ in the FRRF fluorescence (F_o and F_m),
150 however, two extracted chlorophyll measurements from water samples during the
151 survey agree well with F_o (see Figure 8), and here, F_o is used as a proxy for near-
152 surface phytoplankton biomass (e.g., Ellwood et al., 2015).

153 The Moderate Resolution Imaging Spectroradiometer (MODIS, Esaias et al.,
154 1998), launched in 2002 provides satellite-derived estimates of sea surface
155 chlorophyll. Data used here were downloaded on land in near real time from the
156 Ocean Color website (<https://oceancolor.gsfc.nasa.gov/>) and transmitted to the ship
157 once per day.

158 Daily estimates of surface currents were obtained from the AVISO (Archiving,
159 Validation and Interpretation of Satellite Oceanographic data).

160 Least-squares regressions between various quantities were performed using
161 Matlab’s fitlm.m routine, which provides p-values, assuming that the number of
162 degrees of freedom equals the data length minus 2. Linear regression slopes are
163 considered significant if $p < 0.05$.

3. Results

164 3.1 The eddy field off the North Island

165 The satellite-derived surface velocity and chlorophyll fields taken during the
166 cruise on 2 October 2012, just after the spring bloom had started, show several
167 mesoscale (~100 km length scales) eddies, filaments, and other features influencing
168 the SSC field (Figure 2a). Instead of a single Wairarapa eddy, there were at least three
169 distinct anticyclonic eddies (labelled A₁ to A₃) and two distinct cyclonic eddies (C₁,
170 C₂) in the region of the mean Wairarapa eddy.

171 The Rossby number (vorticity/planetary vorticity, $R_o = \zeta / f$) in these eddies
172 typically had a maximum amplitude of 0.15 (Figure 2b), confirming that they were
173 mesoscale, and approximately geostrophic (Mahadevan, 2016). Histograms of R_o for
174 the entire region are slightly skewed positive (positive R_o indicates cyclonic flow in
175 the southern hemisphere), but they become more positively skewed for regions of
176 high SSC (Figure 2c). Moderately high SSC (>1 mg Chl m⁻³) mostly occurred when
177 the absolute value of R_o was less than 0.1, with a preference for cyclonic eddies (
178 Figure 2e). Since the centres of the eddies had R_o about 0.15, this demonstrates that
179 elevated values of SSC typically occurred over the flanks of the eddies rather than at
180 their centres.

181 Conventional theory would suggest that primary production is higher in
182 cyclonic than anticyclonic eddies because mixed layers are shallower in cyclonic
183 eddies (e.g. Doblin et al., 2016), but the satellite image supports the idea that it is the
184 circulation at the edges of the eddies that is important, although not in a simple way.

185 3.2 Physics of Eddy A1

186 The MODIS image of 30 September (Figure 3) showed clear evidence of
187 anticyclonic eddy A1 centred near 180°E, 39°S. Unfortunately, the considerable cloud
188 cover (typical for New Zealand) meant the next clear image was not until 2 October,
189 i.e. after the survey had been started. SSC had evolved into a quite different looking
190 feature by 2 October, and little information can be gleaned on how this change
191 evolved – whether, for example, it was due to the eddy fragmenting and disappearing
192 or whether growth in SSC obscured the eddy.

193 All twelve CTD casts made in the survey had a shallow upper mixed layer
 194 between 15 and 40 m thick sitting over a near-isopycnal layer that extended to
 195 between 250 and 350 m depth (Figure 4). For all casts, except cast 7 (which was the
 196 most inshore station) this deep mixed layer had density, $\sigma_t = 26.499 \pm 0.001 \text{ kg m}^{-3}$
 197 (there was a 0.055 °C and 0.01 range in temperature and salinity in this layer).

198 This near-isopycnal deep layer was almost certainly a remnant mixed layer from
 199 the previous winter mixing, whereas the upper mixed layer reflects the emergence of
 200 new stratification in the spring. Density differences between the upper and remnant
 201 mixed layers ranged between 0.07 and 0.17 kg m^{-3} (Figure 4c). Thus using the
 202 common 0.125 kg m^{-3} density difference criterion for mixed layer depth (e.g., Kara et
 203 al., 2000) would sometimes place the mixed layer as the upper mixed layer and
 204 sometimes as the remnant mixed layer. Here, the upper mixed layer depth (uMLD)
 205 was defined as the depth where the density exceeded the surface value by 0.05 kg m^{-3} ,
 206 and the remnant mixed layer depth (rMLD) was defined as the depth where the
 207 density exceeded the 200m value by 0.05 kg m^{-3} .

208 The upper mixed layer was on average 0.36°C warmer and 0.017 fresher than
 209 the remnant mixed layer. Temperature in the upper mixed layer ranged from 13.58 to
 210 13.93°C, and while the warmest temperature was from noon on 2 October, any effects
 211 of diurnal heating were aliased by spatial variations in the temperature field since the
 212 second-warmest temperature was from 03:30 on 2 October.

213 Gridded upper and remnant mixed layer depths are shown in Figure 5, along
 214 with the velocities at 10-m and 100-m derived from the ADCP. The 100-m velocities
 215 are from near the top of the remnant mixed layer, but are at the depth limit of the
 216 ADCP and are noisier than the 10-m velocities. Velocities from both levels show
 217 anticyclonic circulation. The remnant mixed layer was deeper at the centre of the eddy
 218 (Figure 5b), consistent with anticyclonic eddies, reflecting the fact that these eddies
 219 extend to near the sea floor (Chiswell, 2003). The uMLD was negatively but not
 220 significantly correlated with rMLD (slope = -0.1, $r^2 = 0.24$, p-value = 0.1).

221 The ADCP velocities indicate the anticyclonic eddy was at least ~30 km in
 222 radius with speeds at 10 m depth being ~0.4 m s^{-1} at the northern edge of the survey
 223 (Figure 5). The vorticity ζ calculated from the 10-m velocities was $1.4 \times 10^{-4} \text{ s}^{-1}$
 224 leading to a Rossby number $R_o = \zeta / f$ of -0.16, which puts this eddy into the

225 mesoscale class (i.e. $|R_o| \ll 1$), thus, geostrophy approximately holds, the flow is two-
226 dimensional and the magnitude of the vertical velocity is several orders of magnitude
227 smaller than the horizontal velocities (Mahadevan, 2016).

228 A 3-D view (Figure 6) shows the eddy from a different perspective, and shows
229 the relatively thin warm fresher upper mixed layer sitting over the deeper remnant
230 mixed layer that is warmer and more saline near the centre of the eddy. The remnant
231 mixed layer was deepest near Casts 2 and 10 with shallowest values near the edge of
232 the survey (Casts 6 and 8), whereas the upper mixed layer was shallowest near Cast 2
233 and deepest near Casts 6 and 8.

234 3.3 Chlorophyll distribution within Eddy A1

235 Chlorophyll profiles from the CTD fluorometer (Figure 4d) show NPQ above
236 about 30 m depth in casts made during the day. However, POM was well-mixed in the
237 upper mixed layer in all casts (Figure 4e), likely due to wind-mixing from strong
238 winds on 1 October (see Chiswell et al., 2019). This suggests that all biological
239 quantities (i.e. phytoplankton, dFe, and nutrients) were well mixed in the upper mixed
240 layer, and that near-surface values (in particular, the towed-fish and sea-chest data)
241 well represent upper mixed layer values.

242 The mean one-percent light level during the survey was 60 m, indicating that all
243 upper mixed layers were shallower than the euphotic zone. In every cast, POM was
244 elevated in the upper mixed layer and then decayed with depth to reach values 0.05 to
245 0.09 in the remnant mixed layer. POM in the upper mixed layer (hereafter, POM_u)
246 ranged from 0.57 to 1.0, and visually separates into two groups, which we term high-
247 and low-POM groups, having mean POM_u values of 0.88 and 0.63, respectively
248 (Figure 4e).

249 There is little correlation between POM_u and any physical variable. Inspection
250 of Figure 4 shows there is no immediately obvious relationship between POM group
251 and temperature, salinity, or density. The two warmest, freshest, and lightest upper
252 mixed layers were high-POM, but so too was the coolest, most saline, and densest
253 upper mixed layer. There is also no clear relationship between POM_u and uMLD, for
254 example, two high-POM casts had the shallow uMLD, but other high-POM casts had
255 deep uMLD.

256 Least-squares regressions between POM_u and upper mixed layer temperature or
257 salinity return non-significant results with low correlation-squared ($r^2 < 0.1$, $p > 0.1$, not
258 shown). There is also no statistically significant relationship between POM_u and either
259 uMLD or rMLD ($r^2 \sim 0$, $p > 0.1$, not shown).

260 Much of the reason for these low correlations is because (as is illustrated in the
261 3-D view of the eddy, Figure 6d), POM was high both near the centre of the eddy
262 (casts 2 and 3 with high upper layer T and S, and shallow uMLD) as well as on the
263 flanks of the eddy (casts 6 to 8 with low upper layer T and S, and deeper uMLD).

264 Figure 7 shows the macronutrient concentrations (NH_4 , NO_3+NO_2 , PO_4 , SiO_4)
265 as a function of depth, colour-coded by POM group. Unfortunately, the depth levels
266 for the nutrient sampling were pre-determined before the survey, and the 30-m level
267 falls near the base of the upper mixed layer in many casts (see Figure 4), which
268 complicates interpretation (i.e. not all these 30-m nutrient values were unambiguously
269 from the upper mixed layer). There is considerable scatter, but most macronutrients
270 increased with depth. At 10 m depth (which is the only level consistently in the upper
271 mixed layer), NO_3+NO_2 and PO_4 showed lower values for the casts that had high
272 POM_u , which would be consistent with the uptake of these nutrients by
273 phytoplankton, but SiO_4 and NH_4 show no evidence of uptake.

274 While Chl and POM_u from the CTD casts, and F_o from the FRRF are all proxies
275 for phytoplankton biomass, F_o is the more useful proxy to compare with dFe because
276 of its more frequent sampling, and it is worth demonstrating that F_o correlates well
277 with the other measures of phytoplankton biomass. Surface chlorophyll from the
278 night-time CTD casts and POM_u from all casts align well with F_o (r^2 values of 0.74
279 and 0.38, and p-values of 0.007 and 0.033, respectively, Figure 8a). Similarly, F_o and
280 POM_u both compare well with SSC from the 2 October MODIS image (which was
281 taken about the time cast 4 was made). There are some discrepancies, but except for
282 the last two casts (by which time the satellite image was 12 to 16 hours old), high F_o
283 and POM_u generally overlie regions of high SSC, and low F_o and POM_u overlie
284 regions of low SSC (Figure 8b).

285 Figure 9a presents F_o along with dFe from the tow fish. The ship passed through
286 a region of elevated phytoplankton biomass during the morning of 2 October,
287 followed by low biomass in the afternoon and then elevated biomass during the night

288 (these are the regions near the centre and on the flanks of the eddy seen in Figure 6).
 289 A least-square regression (with F_0 values interpolated to the dFe sample times)
 290 suggests that dFe is positively correlated with F_0 ($r^2 = 0.27$; $p = 0.006$, Figure 9d).
 291 That dFe is correlated with phytoplankton biomass is also illustrated when dFe is
 292 superimposed on satellite SSC where high values of dFe coincide with high SSC
 293 (Figure 8b).

294 Dissolved iron, dFe, is correlated with uMLD ($r^2 = 0.18$; $p = 0.03$, Figure 9b, e),
 295 although since the operational procedure was to deploy the trace-metal tow fish
 296 between the CTD casts, the dFe values were taken in slightly different locations, and
 297 there is the possibility that small-scale patchiness confounds this relationship. Because
 298 dFe increases with depth (Chiswell et al., 2019, Figure 4), this suggests that part of the
 299 dFe signal could be due to entrainment of higher-dFe water as the upper mixed level
 300 deepened.

301 The upper mixed layer depth and F_0 were uncorrelated ($r^2 = 0.01$; $p = 0.175$, not
 302 shown), so they can be used as independent variables in a linear predictor to estimate
 303 dFe,

$$304 \quad dFe_{est} = a \times uMLD + b \times F_0 + c .$$

305 A multiple least-squares linear regression using the data shown in Figure 9, returns a
 306 $= 0.003 \pm 0.0018 \text{ nmol L}^{-1} \text{ m}^{-1}$ and $b = 0.37 \pm 0.14 \text{ nmol L}^{-1}$, and $c = -0.18 \pm 0.10 \text{ nmol}$
 307 L^{-1} , with an overall p-value of 0.005. A fit with uMLD and F_0 de-meaned and
 308 normalised by their respective standard deviations, returns $a = 0.019 \pm 0.01$, $b = 0.$
 309 025 ± 0.1 , and $c = 0.17 \pm 0.1$, indicating that the contribution to iron uptake/loss by
 310 normalised fluorescence variations is about 1.3 times that of normalised mixed layer
 311 depth variations. Whether computed dimensionally or non-dimensionally, the fit,
 312 dFe_{est} , accounts for 36% of the variance in dFe (Figure 9f). The implications of this
 313 fit are discussed in detail later, but for now we simply comment that positive a is
 314 consistent with entrainment of high-dFe water from depth, whereas positive b is
 315 unexpected since one might expect uptake of iron to lead to negative b .

4. Discussion

316 The remnant mixed layer in eleven of the twelve casts had near-identical density
 317 ($\sigma_t = 26.499 \pm 0.001 \text{ kg m}^{-3}$), suggesting that the deeper water within Eddy A₁ had a

318 common origin. This eddy was likely formed near East Cape where one to two eddies
319 are shed per year that propagate south-west, but are topographically blocked by the
320 Chatham Rise so that in the mean, the circulation is as shown in Figure 1 (Chiswell,
321 2005). The CTD casts were made only to 500 m at most, so that the full water-column
322 baroclinic structure of eddy A₁ cannot be determined from this survey. However, the
323 mean Wairarapa Eddy extends to at least 1000 m depth (Roemmich and Sutton,
324 1998), and individual mesoscale eddies are as deep (Chiswell, 2003), hence the
325 observation that the remnant mixed layer structure is consistent with anticyclonic
326 rotation (i.e. deeper mixed layer in the centre of the eddy, Figure 5). The shallow
327 upper mixed layer (mean depth = 30 m) had negligible impact on the circulation.

328 The upper mixed layer during the survey was on average 0.37°C warmer and
329 0.019 fresher than the remnant mixed layer. This temperature difference is similar to
330 that seen earlier in the cruise as the ocean became warmer and began to stratify
331 (Chiswell et al., 2019). The decrease in salinity in the upper mixed layer can be
332 accounted for by local precipitation exceeding evaporation (0.01m of rain would lead
333 to about 0.01 salinity decrease if mixed over a 30-m upper mixed layer). Thus, the T-
334 S properties of the upper mixed layer were likely to have been heavily modified by
335 recent heating and rainfall, and so cannot be used to determine if regions of high- and
336 low- POM were of coastal vs offshore origin (or vice-versa).

337 In every cast, the upper mixed layer showed more POM than in the remnant
338 mixed layer (Figure 4e), indicating more phytoplankton biomass in the upper mixed
339 layer. A complete analysis of this production is beyond the scope of this article,
340 however, higher phytoplankton biomass in the upper mixed layer is broadly consistent
341 with the Onset of Stratification model for spring blooms (Chiswell, 2011), but the lack
342 of correlation between POM_u or F_o with either depth or temperature of the upper
343 mixed layer suggests that the amount of phytoplankton biomass in the upper mixed
344 layer cannot be ascribed to the age (assuming that a cooler upper mixed layer is
345 formed more recently than a warmer upper mixed layer), or depth of the upper mixed
346 layer. In other words, it seems unlikely that regions of low-biomass had stratified
347 more recently or were earlier in the spring bloom cycle than regions of high-biomass.

348 Apart from a suggestion of nitrate and phosphate uptake, there was no
349 statistically significant relationship between phytoplankton biomass in the upper

350 mixed layer and macronutrients, although this lack of relationship may well have been
351 due more to the lack of vertical nutrient sampling.

352 There was also no statistically significant relationship between upper mixed
353 layer chlorophyll or POM and the remnant mixed layer depth. At first glance, this
354 may not be surprising since it is unlikely that a mixed layer some 200-300 m below
355 the upper mixed layer has a role in determining the upper mixed layer production.
356 However, it has been suggested (Uchida et al., 2020) that vertical velocities associated
357 with mesoscale eddies can bring iron-rich water from depth into the mixed layer
358 where it can be consumed by phytoplankton, and their model simulations show much
359 higher dFe in eddy-rich areas. Crucially, however, within their eddies they found a
360 negative relationship between dFe and phytoplankton biomass, which we did not see,
361 so while we cannot exclude such a mechanism from playing a role, it seems unlikely
362 that it is the dominant mechanism driving the production within the eddy field in our
363 study.

364 The only significant relationships we could find here were between dFe and
365 chlorophyll or POM, and between dFe and upper mixed layer depth, with a simple
366 linear model accounting for 36% of the variance in dFe.

367 While dFe was not measured within the remnant mixed layer during this survey,
368 three days earlier in the cruise it was about 0.35 nmol L⁻¹ (see Figure 4 in Chiswell et
369 al., 2019). About 18% of the near-surface dFe variance can be explained by deepening
370 of the upper mixed layer, and to the extent that the uML deepens through vertical
371 mixing, this suggests entrainment of high-dFe water from the remnant mixed layer
372 into the upper mixed layer (Figure 9e).

373 About 27% of the variance in near-surface dFe is explained by increased F_o ,
374 (Figure 9d). This positive relationship between dFe and phytoplankton biomass is a
375 little surprising because one would expect uptake during a bloom to lead to a negative
376 relationship. When combined with the satellite images showing apparent lateral
377 entrainment of neritic water, we suggest that the most likely explanation for this result
378 is that the eddies transport high-dFe coastal water offshore. This conclusion is
379 supported by results from the same region in 2008, where Ellwood et al. (2014) found
380 that mixed layer Mn:Fe and Mn:Al ratios were elevated compared to crustal values,

381 and based on these ratios and particle-backtracking simulations, suggested the most
382 likely source of iron was continental water.

383 Even so, if the coastal water had uniform iron concentration, uptake would lead
384 to a negative relationship between dFe and phytoplankton biomass. There are two
385 potential reasons for a positive relationship. One is that our results are confounded by
386 atmospheric deposition, the other is that coastal water has such a large range in
387 dissolved iron that uptake is obscured by this variability.

388 If phytoplankton are iron-limited, patchy atmospheric deposition of iron could
389 lead to patches of phytoplankton, and thus to a positive relationship between dFe and
390 phytoplankton, although it is by no means certain that deposition leads to increased
391 dFe since lithogenic particles associated with atmospheric deposition can remove
392 soluble iron in the water column via scavenging (Tagliabue et al., 2017).

393 Atmospheric deposition of dust in the New Zealand region has long been
394 ascribed to Australian dust storms (Boyd et al., 2004; Kidson, 1930; Mahowald et al.,
395 2009), but there are not sufficient data to determine the temporal and spatial scales of
396 the deposition in the region. Ellwood et al. (2018; their Figure 5) present a map of
397 annual mean Fe deposition based on modelling of global dust distributions (Albani et
398 al., 2014) that shows broad area of deposition extending east of Australia and
399 covering New Zealand. This modelled deposition has little of the spatial structure of
400 surface chlorophyll seen in Figure 1, in particular there is no evidence of high
401 deposition along the STF, and since the STF is considered to be region of high dFe
402 (Banse and English, 1997; Boyd et al., 2004) this argues that deposition is not the
403 major source of iron to the STF. Such featureless deposition might be expected in the
404 annual mean, but various attempts to investigate the role of atmospheric deposition
405 from satellite imagery have not provided more precise information on the temporal
406 and spatial scales of deposition at shorter time scales. Boyd et al. (2004) inferred
407 oceanic supply of dFe from episodic increases in chlorophyll concentrations in SAW
408 seen in ocean colour images between 1997 and 2001, and found no evidence that
409 these events were mediated by atmospheric iron supply, although they also could not
410 explain these events from lateral advection or vertical mixing, and did comment that
411 dust storms during this time sent plumes over both STW and SAW. They concluded
412 that more data are needed, including on rainfall patterns in relation to dust plume
413 trajectories to distinguish wet from dry deposition.

414 With little to no measurements of atmospheric deposition of iron in the region,
415 the role of deposition in this mesoscale eddy system must remain an open question.

416 This then raises the question of whether our implied finding that dFe is high and
417 variable in coastal water is consistent with previous work. There is surprisingly little
418 information on dFe in coastal North Island water, however, it is reasonable to assume
419 that coastal water, in general is higher in dFe than offshore water. For example,
420 Hutchins et al. (1998) found dFe in California coastal water to vary between <0.1 and
421 >8.0 nmol L^{-1} , and explained this high variability was due to uneven distributions of
422 sources of iron such as rivers and resuspension of shelf sediments. More locally,
423 Sander et al. (2015) found dFe on average to be 1.0 ± 0.4 nmol L^{-1} in neritic water on
424 the Otago shelf, compared to 0.2 ± 0.1 nmol L^{-1} offshore in SAW. Croot and Hunter
425 (1998) reported dFe as high as 6 nmol L^{-1} on the Otago shelf, which they suggested
426 was due to wind-induced upwelling rather than fluvial input. Upwelling occurs around
427 the North Island east coast (Sharples and Greig, 1998) and may also be a significant
428 source of dFe. On balance, it seems that high and variable dFe in coastal water is a
429 reasonable finding.

430 Evidence that dFe is limiting in south-west Pacific Ocean STW is more scarce,
431 perhaps because STW is generally considered oligotrophic in macronutrients and
432 there has been relatively little interest in iron in these waters. However, Ellwood et al.
433 (2008) found winter dFe levels ~ 0.1 nmol L^{-1} within and across the Subtropical Front
434 in the Tasman Sea. Near 30°S , Ellwood et al. (2018) found surface dFe less than 0.2
435 nmol L^{-1} in the central Tasman Sea. Furthermore, phytoplankton located north of the
436 Tasman Front (i.e. in STW, but to the west of our study region) were considered to be
437 near the threshold for iron limitation (Ellwood et al., 2013).

438 In summary, the main results of this study are that the mesoscale eddies off the
439 east coast of New Zealand are deep, with their circulation driven by the deep
440 baroclinic structure, whereas (at least during the spring bloom) the surface production
441 is largely constrained to a shallow upper mixed layer. From a biological perspective,
442 the main role of the mesoscale circulation is to mix high-dFe neritic water with low-
443 dFe offshore STW. Since extensive mixing of the two water masses extends several
444 hundred km offshore, we conclude that a significant component of production within
445 the STF is likely determined by mesoscale eddy induced zonal advection of iron. This
446 conclusion supports those made by Graham et al. (2015), who suggested that

447 bioavailable iron from the continental shelves is entrained into western boundary
448 currents and then advected along the STF.

449

Acknowledgements

450 We thank all those involved during the cruise, including the Master and crew of
451 RV Tangaroa for their contribution to the data collection. Simon Wood provided daily
452 satellite images to aid us in locating the eddy. The MODIS ocean colour data are
453 made freely available by NASA's Ocean Color Program supported by the Ocean
454 Biology Processing Group (OBPG) at NASA's Goddard Space Flight Center
455 (<https://oceancolor.gsfc.nasa.gov/>). AVISO ocean altimeter data are made freely
456 available and distributed by AVISO (<https://www.aviso.altimetry.fr/>). This work was
457 funded by the New Zealand Ministry of Business, Innovation and Employment
458 through several grants to NIWA. MJE was supported by the Australian Research
459 Council [grants DP170102108 and DP130100679]. AM was supported by the Natural
460 Environmental Research Council [grant NERC NE/H004475/1 awarded to Maeve
461 Lohan]. The IAEA is grateful to the Government of the Principality of Monaco for the
462 support provided to its Environment Laboratories. We also thank the reviewers of this
463 article for their conscientious reviews.

References

464

Figures

465

466 Figure 1. Mean sea surface chlorophyll (SSC) from 2002 - 2016 derived from the
 467 Moderate Resolution Imaging Spectroradiometer (MODIS) on NASA's
 468 Aqua satellite. **a)** Mean SSC for the southern hemisphere; and **b)** Mean
 469 SSC for the New Zealand region. Black vectors are mean sea surface
 470 velocity derived from AVISO ocean altimeter data. Red lines are 0.5 and
 471 0.75 mg Chl m⁻³ contour levels, dash-dot black line is 1000 m isobath,
 472 showing the Chatham Rise extending east of New Zealand. EC, WE, and
 473 CI indicate East Cape, Wairarapa Eddy, and Chatham Islands.

474

475 Figure 2. **a)** Sea surface chlorophyll (SSC) from 2 October 2012 derived from the
 476 Moderate Resolution Imaging Spectroradiometer (MODIS) on NASA's
 477 Aqua satellite. Vectors are surface currents derived from AVISO ocean
 478 altimeter data. Labels A, and C indicate anti-cyclonic and cyclonic eddies,
 479 numbered eddies are referred to in the text. The blue line near eddy A₁
 480 indicates the cruise track during the one-day survey; **b)** Rossby number
 481 (vorticity divided by planetary vorticity, $R_0 = \zeta / f$) for 2 October 2012.
 482 Dashed contours indicate $R_0 = \pm 0.1$; and **c-f)** Histograms of Rossby
 483 number for regions where SSC exceeds various levels. Dash-dot line in **a**
 484 and **b** indicates the 250 m isobath, and data are only shown where the water
 485 depth is greater than 250 m to exclude the coastal zone.

486

487 Figure 3. Moderate Resolution Imaging Spectroradiometer (MODIS) images of sea
 488 surface chlorophyll (SSC) from 30 September to 4 October, 2012. **a)** 30
 489 September; **b)** 2 October; **c)** 4 October; and **d)** enlarged view of 2 October,
 490 along with upper mixed-layer velocity from the ship's ADCP (red vectors).
 491 Numbered circles show the CTD cast locations from the one-day survey
 492 made in the anticyclonic eddy A1 shown in Figure 2. Colours of the circles
 493 in **d)** indicate whether the casts were considered high-POM (green) or low-
 494 POM (blue, see text). Black vectors are surface currents derived from
 495 AVISO satellite altimetry.

496

497 Figure 4. Profiles from the 12 CTD casts made during the one-day survey. **a)**
 498 Temperature (T); **b)** Salinity (S); **c)** Density (σ_t); **d)** Chlorophyll (Chl); **e)**
 499 Particulate organic matter (POM); **f)** Temperature vs. salinity (T-S); and **g)**
 500 POM vs Chl for the upper 5 to 30 m from CTD casts made over the entire
 501 cruise. Casts have been colour-coded by two groups, high-POM (green) and
 502 low-POM (blue). Horizontal blue line shows the mean one-percent light
 503 level (Z_{100}) from the daylight casts. Horizontal red dashed lines show the
 504 nutrient sample depths. Note the change of depth scale between upper and
 505 lower panels. Magenta lines in plots are from cast 7, which was closest
 506 inshore and had lowest temperature and salinity in the remnant mixed layer.

507

508 Figure 5. Gridded mixed-layer depths derived from CTD casts shown in Figure 4.
 509 **a)** Upper mixed layer depth (uMLD); and **b)** Remnant mixed layer depth
 510 (rMLD). Red and blue vectors are ADCP-derived velocities at 10 m and
 511 100 depths, respectively, showing anticyclonic circulation.

512

513 Figure 6. Three-dimensional views of **a)** Temperature (T); **b)** Salinity (S); **c)**
 514 Density (σ_t); and **d)** Particulate organic matter (POM) from the one-day
 515 CTD survey. Red vectors are ADCP-derived velocities at 10 m showing
 516 anticyclonic circulation. Vertical lines indicate upper mixed-layer depths
 517 and are colour-coded as high POM_u (green) and low POM_u (blue). Black
 518 and white dashed lines indicate the depths of the remnant and upper mixed
 519 layers, respectively. Numbers in **d)** are the cast number. Note change of
 520 vertical (depth) scale between **a)**, **b)** and **c)**, **d)**.

521

522 Figure 7. Macronutrient profiles, from the CTD survey. **a)** Ammonium (NH_4); **b)**
 523 Nitrate plus nitrite (NO_3+NO_2); **c)** Phosphate (PO_4); and **d)** Silicate (SiO_4).
 524 Values have been colour-coded as high-POM (green) and low-POM (blue)
 525 profiles (see Figure 4). Error bars are one standard deviation of analytical
 526 errors and do not include CTD sampling errors.

527

528 Figure 8. **a)** Time series of fluorescence (F_o) from Fast Repetition Rate
 529 Fluorometer (green line, and 1-hour smoothed, dark-green line), surface
 530 chlorophyll from night-time CTD casts (dark green squares), extracted
 531 chlorophyll from water samples at 10 m (red diamonds), and upper-mixed
 532 layer particulate organic matter (POM_u) from CTD casts (circles). POM_u
 533 has been colour-coded as high- POM_u (green) and low- POM_u (blue); and
 534 **b)** Scatter plots of F_o , POM_u , and dFe superimposed over sea surface
 535 chlorophyll (SSC) derived from Moderate Resolution Imaging
 536 Spectroradiometer for 2 October 2012 (see Figure 3). Note the colour scale
 537 for SSC is as shown in Figure 3. Numbers in circles indicate CTD cast
 538 number. Casts 2 to 5 were made during daylight.

539

540 Figure 9. Dissolved iron from underway tow fish (dFe), fluorescence from Fast
 541 Repetition Rate fluorometer (F_o), upper mixed layer depth from CTD casts
 542 (uMLD), and fit of dFe from linear model ($dFe_{est} = a \times uMLD + b \times F_o + c$).
 543 **a)** dFe and F_o ; **b)** dFe and uMLD; **c)** dFe and dFe_{est} ; and **d-f)** linear
 544 regressions of F_o , uMLD, and dFe_{est} against dFe , where F_o and uMLD, has
 545 been interpolated to dFe sample times. All slopes are considered significant
 546 ($p < .05$). In **a)** and **b)** scales for F_o and uMLD are shown to the right of the
 547 axes.

548

549 Abraham, E.R., Law, C.S., Boyd, P.W., Lavender, S.J., Maldonado, M.T., Bowie,
 550 A.R., 2000. Importance of stirring in the development of an iron-fertilized
 551 phytoplankton bloom. *Nature* 407 (6805), 727-730. <https://doi.org/10.1038/35037555>.
 552 Albani, S., Mahowald, N.M., Perry, A.T., Scanza, R.A., Zender, C.S., Heavens, N.G.,
 553 Maggi, V., Kok, J.F., Otto-Bliesner, B.L., 2014. Improved dust representation in the
 554 Community Atmosphere Model. *Journal of Advances in Modeling Earth Systems* 6
 555 (3), 541-570. <https://doi.org/10.1002/2013MS000279>.
 556 Banse, K., English, D.C., 1997. Near-surface phytoplankton pigment from the Coastal
 557 Zone Color Scanner in the Subantarctic region southeast of New Zealand. *Marine*
 558 *Ecology Progress Series* 156, 51-66. <https://doi.org/10.3354/meps156051>.
 559 Bishop, J.K.B., 1999. Transmissometer measurement of POC. *Deep Sea Research*
 560 *Part I: Oceanographic Research Papers* 46 (2), 353-369.
 561 [https://doi.org/10.1016/S0967-0637\(98\)00069-7](https://doi.org/10.1016/S0967-0637(98)00069-7).

- 562 Blain, S., Bonnet, S., Guieu, C., 2008. Dissolved iron distribution in the tropical and
563 sub tropical South Eastern Pacific. *Biogeosciences* 5 (1), 269-280.
564 <https://doi.org/10.5194/bg-5-269-2008>.
- 565 Boyd, P., LaRoche, J., Gall, M., Frew, R., McKay, R.M.L., 1999. Role of iron, light,
566 and silicate in controlling algal biomass in subantarctic waters SE of New Zealand.
567 *Journal of Geophysical Research: Oceans* 104 (C6), 13395-13408.
568 <https://doi.org/10.1029/1999JC900009>.
- 569 Boyd, P.W., McTainsh, G., Sherlock, V., Richardson, K., Nichol, S., Ellwood, M.,
570 Frew, R., 2004. Episodic enhancement of phytoplankton stocks in New Zealand
571 subantarctic waters: Contribution of atmospheric and oceanic iron supply. *Global*
572 *Biogeochemical Cycles* 18 (1). <https://doi.org/10.1029/2002GB002020>.
- 573 Boyd, P.W., Strzepek, R., Chiswell, S., Chang, H., DeBruyn, J.M., Ellwood, M.,
574 Keenan, S., King, A.L., Maas, E.W., Nodder, S., Sander, S.G., Sutton, P., Twining,
575 B.S., Wilhelm, S.W., Hutchins, D.A., 2012. Microbial control of diatom bloom
576 dynamics in the open ocean. *Geophysical Research Letters* 39 (18), L18601.
577 <https://doi.org/10.1029/2012gl053448>.
- 578 Boyd, P.W., Watson, A.J., Law, C.S., Abraham, E.R., Trull, T., Murdoch, R., Bakker,
579 D.C., Bowie, A.R., Buesseler, K.O., Chang, H., Charette, M., Croot, P., Downing, K.,
580 Frew, R., Gall, M., Hadfield, M., Hall, J., Harvey, M., Jameson, G., LaRoche, J.,
581 Liddicoat, M., Ling, R., Maldonado, M.T., McKay, R.M., Nodder, S., Pickmere, S.,
582 Pridmore, R., Rintoul, S., Safi, K., Sutton, P., Strzepek, R., Tanneberger, K., Turner,
583 S., Waite, A., Zeldis, J., 2000. A mesoscale phytoplankton bloom in the polar
584 Southern Ocean stimulated by iron fertilization. *Nature* 407 (6805), 695-702.
585 <https://doi.org/10.1038/35037500>.
- 586 Bradford, J.M., Heath, R.A., Chang, F.H., Hay, C.H., 1982. The effect of warm-core
587 eddies on oceanic productivity off northeastern New Zealand. *Deep Sea Research Part*
588 *A. Oceanographic Research Papers* 29 (12), 1501-1516. [https://doi.org/10.1016/0198-](https://doi.org/10.1016/0198-0149(82)90039-5)
589 [0149\(82\)90039-5](https://doi.org/10.1016/0198-0149(82)90039-5).
- 590 Carberry, L., Roesler, C., Drapeau, S., 2019. Correcting in situ chlorophyll
591 fluorescence time-series observations for nonphotochemical quenching and tidal
592 variability reveals nonconservative phytoplankton variability in coastal waters.
593 *Limnology and Oceanography: Methods* 17 (8), 462-473.
594 <https://doi.org/10.1002/lom3.10325>.
- 595 Chiswell, S.M., 2003. Circulation within the Wairarapa Eddy, New Zealand. *New*
596 *Zealand Journal of Marine and Freshwater Research* 37, 691-704.
597 <https://doi.org/10.1080/00288330.2003.9517199>.
- 598 Chiswell, S.M., 2005. Mean and variability in the Wairarapa and Hikurangi Eddies,
599 New Zealand. *New Zealand Journal of Marine and Freshwater Research* 39 (1), 121-
600 134. <https://doi.org/10.1080/00288330.2005.9517295>.
- 601 Chiswell, S.M., 2011. Annual cycles and spring blooms in phytoplankton: don't
602 abandon Sverdrup completely. *Marine Ecology Progress Series* 443, 39-50.
603 <https://doi.org/10.3354/meps09453>.
- 604 Chiswell, S.M., Gutiérrez-Rodríguez, A., Gall, M., Safi, K., Strzepek, R., Decima, M.,
605 Nodder, S.D., 2022. Phytoplankton Phenology and Net Primary Production from
606 Biogeochemical Argo Floats in the South-West Pacific Ocean. *Deep Sea Research*
607 submitted.
- 608 Chiswell, S.M., Safi, K.A., Sander, S.G., Strzepek, R., Ellwood, M.J., Milne, A.,
609 Boyd, P.W., 2019. Exploring mechanisms for spring bloom evolution: contrasting
610 2008 and 2012 blooms in the southwest Pacific Ocean. *Journal of Plankton Research*.
611 <https://doi.org/10.1093/plankt/fbz017>.

- 612 Croot, P.L., Hunter, K.A., 1998. Trace metal distributions across the continental shelf
613 near Otago Peninsula, New Zealand. *Marine Chemistry* 62 (3-4), 185-201.
614 [https://doi.org/10.1016/S0304-4203\(98\)00036-X](https://doi.org/10.1016/S0304-4203(98)00036-X).
- 615 Doblin, M.A., Petrou, K., Sinutok, S., Seymour, J.R., Messer, L.F., Brown, M.V.,
616 Norman, L., Everett, J.D., McInnes, A.S., Ralph, P.J., Thompson, P.A., Hassler, C.S.,
617 2016. Nutrient uplift in a cyclonic eddy increases diversity, primary productivity and
618 iron demand of microbial communities relative to a western boundary current. *PeerJ*
619 4, e1973-e1973. <https://doi.org/10.7717/peerj.1973>.
- 620 Ellwood, M.J., Bowie, A.R., Baker, A., Gault-Ringold, M., Hassler, C., Law, C.S.,
621 Maher, W.A., Marriner, A., Nodder, S., Sander, S., Stevens, C., Townsend, A., van
622 der Merwe, P., Woodward, E.M.S., Wuttig, K., Boyd, P.W., 2018. Insights Into the
623 Biogeochemical Cycling of Iron, Nitrate, and Phosphate Across a 5,300 km South
624 Pacific Zonal Section (153°E-150°W). *Global Biogeochemical Cycles* 32 (2), 187-
625 207. <https://doi.org/10.1002/2017gb005736>.
- 626 Ellwood, M.J., Boyd, P.W., Sutton, P., 2008. Winter-time dissolved iron and nutrient
627 distributions in the Subantarctic Zone from 40–52S; 155–160E. *Geophysical Research*
628 *Letters* 35 (11). <https://doi.org/10.1029/2008gl033699>.
- 629 Ellwood, M.J., Hutchins, D.A., Lohan, M.C., Milne, A., Nasemann, P., Nodder, S.D.,
630 Sander, S.G., Strzepek, R., Wilhelm, S.W., Boyd, P.W., 2015. Iron stable isotopes
631 track pelagic iron cycling during a subtropical phytoplankton bloom. *Proceedings of*
632 *the National Academy of Sciences of the United States of America* 112 (1), E15-E20.
633 <https://doi.org/10.1073/pnas.1421576112>.
- 634 Ellwood, M.J., Law, C.S., Hall, J., Woodward, E.M.S., Strzepek, R., Kuparinen, J.,
635 Thompson, K., Pickmere, S., Sutton, P., Boyd, P.W., 2013. Relationships between
636 nutrient stocks and inventories and phytoplankton physiological status along an
637 oligotrophic meridional transect in the Tasman Sea. *Deep Sea Research Part I:*
638 *Oceanographic Research Papers* 72, 102-120.
639 <https://doi.org/10.1016/j.dsr.2012.11.001>.
- 640 Ellwood, M.J., Nodder, S.D., King, A.L., Hutchins, D.A., Wilhelm, S.W., Boyd,
641 P.W., 2014. Pelagic iron cycling during the subtropical spring bloom, east of New
642 Zealand. *Marine Chemistry* 160, 18-33.
643 <https://doi.org/10.1016/j.marchem.2014.01.004>.
- 644 Esaias, W.E., Abbott, M.R., Barton, I., Brown, O.B., Campbell, J.W., Carder, K.L.,
645 Clark, D.K., Evans, R.H., Hoge, F.E., Gordon, H.R., Balch, W.M., Letelier, R.,
646 Minnett, P.J., 1998. An overview of MODIS capabilities for ocean science
647 observations. *IEEE Transactions on Geoscience and Remote Sensing* 36 (4), 1250-
648 1265. <https://doi.org/10.1109/36.701076>.
- 649 Floor, G.H., Clough, R., Lohan, M.C., Ussher, S.J., Worsfold, P.J., Quetel, C.R.,
650 2015. Combined uncertainty estimation for the determination of the dissolved iron
651 amount content in seawater using flow injection with chemiluminescence detection.
652 *Limnology and Oceanography Methods* 13 (12), 673-686.
653 <https://doi.org/10.1002/lom3.10057>.
- 654 Graham, R.M., De Boer, A.M., van Sebille, E., Kohfeld, K.E., Schlosser, C., 2015.
655 Inferring source regions and supply mechanisms of iron in the Southern Ocean from
656 satellite chlorophyll data. *Deep Sea Research Part I: Oceanographic Research Papers*
657 104 (Supplement C), 9-25. <https://doi.org/10.1016/j.dsr.2015.05.007>.
- 658 Hutchins, D.A., DiTullio, G.R., Zhang, Y., Bruland, K.W., 1998. An iron limitation
659 mosaic in the California upwelling regime. *Limnology and Oceanography* 43 (6),
660 1037-1054. <https://doi.org/10.4319/lo.1998.43.6.1037>.

- 661 Jickells, T., Moore, C.M., 2015. The importance of Atmospheric Deposition for
662 Ocean Productivity. *Annual Review of Ecology, Evolution, and Systematics* 46 (1),
663 481-501. <https://doi.org/10.1146/annurev-ecolsys-112414-054118>.
- 664 Johnson, K.S., Plant, J.N., Dunne, J.P., Talley, L.D., Sarmiento, J.L., 2017. Annual
665 nitrate drawdown observed by SOCCOM profiling floats and the relationship to
666 annual net community production. *Journal of Geophysical Research: Oceans* 122 (8),
667 6668-6683. <https://doi.org/10.1002/2017JC012839>.
- 668 Kara, A.B., Rochford, P.A., Hurlburt, H.E., 2000. An optimal definition for ocean
669 mixed layer depth. *Journal of Geophysical Research: Oceans* 105 (C7), 16803-16821.
670 <https://doi.org/10.1029/2000JC900072>.
- 671 Kidson, E., 1930. Australian Origin of Red Rain in New Zealand. *Nature* 125 (3150),
672 410-410. <https://doi.org/10.1038/125410a0>.
- 673 Mahadevan, A., 2016. The Impact of Submesoscale Physics on Primary Productivity
674 of Plankton. *Ann Rev Mar Sci* 8, 161-184. <https://doi.org/10.1146/annurev-marine-010814-015912>.
- 676 Mahowald, N.M., Engelstaedter, S., Luo, C., Sealy, A., Artaxo, P., Benitez-Nelson,
677 C., Bonnet, S., Chen, Y., Chuang, P.Y., Cohen, D.D., Dulac, F., Herut, B., Johansen,
678 A.M., Kubilay, N., Losno, R., Maenhaut, W., Paytan, A., Prospero, J.M., Shank,
679 L.M., Siefert, R.L., 2009. Atmospheric iron deposition: global distribution,
680 variability, and human perturbations. *Ann Rev Mar Sci* 1, 245-278.
681 <https://doi.org/10.1146/annurev.marine.010908.163727>.
- 682 Obata, H., Karatani, H., Nakayama, E., 2002. Automated determination of iron in
683 seawater by chelating resin concentration and chemiluminescence detection.
684 *Analytical Chemistry* 65 (11), 1524-1528. <https://doi.org/10.1021/ac00059a007>.
- 685 Pickmere, S.E., 1998. Biological effects of cross-shelf water transfer programme
686 nutrient report. NIWA Internal Report, p. 5.
- 687 Roemmich, D., Sutton, P., 1998. The mean and variability of ocean circulation past
688 northern New Zealand: Determining the representativeness of hydrographic
689 climatologies. *Journal of Geophysical Research: Oceans* 103 (C6), 13041-13054.
690 <https://doi.org/10.1029/98jc00583>.
- 691 Sander, S.G., Tian, F., Ibisami, E.B., Currie, K.I., Hunter, K.A., Frew, R.D., 2015.
692 Spatial and seasonal variations of iron speciation in surface waters of the Subantarctic
693 front and the Otago Continental Shelf. *Marine Chemistry* 173, 114-124.
694 <https://doi.org/10.1016/j.marchem.2014.09.001>.
- 695 Sharples, J., Greig, M.J.N., 1998. Tidal currents, mean flows, and upwelling on the
696 north-east shelf of New Zealand. *New Zealand Journal of Marine and Freshwater
697 Research* 32 (2), 215-231. <https://doi.org/10.1080/00288330.1998.9516821>.
- 698 Tagliabue, A., Bowie, A.R., Boyd, P.W., Buck, K.N., Johnson, K.S., Saito, M.A.,
699 2017. The integral role of iron in ocean biogeochemistry. *Nature* 543 (7643), 51-59.
700 <https://doi.org/10.1038/nature21058>.
- 701 Uchida, T., Balwada, D., P. Abernathy, R., A. McKinley, G., K. Smith, S., Lévy, M.,
702 2020. Vertical eddy iron fluxes support primary production in the open Southern
703 Ocean. *Nature Communications* 11 (1), 1125. <https://doi.org/10.1038/s41467-020-14955-0>.
- 704
- 705 Waite, A.M., Pesant, S., Griffin, D.A., Thompson, P.A., Holl, C.M., 2007.
706 Oceanography, primary production and dissolved inorganic nitrogen uptake in two
707 Leeuwin Current eddies. *Deep Sea Research Part II: Topical Studies in Oceanography*
708 54 (8), 981-1002. <https://doi.org/10.1016/j.dsr2.2007.03.001>.
- 709